


State of Oregon
Department of Geology and Mineral Industries
Vicki S. McConnell, State Geologist

Open-File Report O-07-05

GEOLOGIC COMPILATION MAP OF PART OF THE UPPER KLAMATH BASIN, KLAMATH COUNTY, OREGON

Compiled by
Margaret D. Jenks¹



2007

¹Oregon Department of Geology and Mineral Industries, 800 NE Oregon St., Suite 965, Portland, Oregon 97232

NOTICE

No warranty, expressed or implied, is made regarding the accuracy or utility of the data described and/or contained herein, nor shall the act of distribution constitute any such warranty. This disclaimer applies both to individual use of the data and aggregate use with other data. The Oregon Department of Geology and Mineral Industries shall not be held liable for improper or incorrect use of these data. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

Oregon Department of Geology and Mineral Industries Open-File Report O-07-05
Published in conformance with ORS 516.030

For copies of this publication or other information about Oregon's geology and natural resources, contact:

Nature of the Northwest Information Center
800 NE Oregon Street #5
Portland, Oregon 97232
(503) 872-2750
<http://www.naturenw.org>

or these DOGAMI field offices:

Baker City Field Office
1510 Campbell St.
Baker City, OR 97814-3442
Telephone (541) 523-3133
Fax (541) 523-5992

Grants Pass Field Office
5375 Monument Drive
Grants Pass, OR 97526
Telephone (541) 476-2496
Fax (541) 474-3158

For additional information:
Administrative Offices
800 NE Oregon Street, Suite 965
Portland, OR 97232
Telephone (971) 673-1555
Fax (971) 673-1562
<http://www.oregongeology.com>
<http://egov.oregon.gov/DOGAMI/>

TABLE OF CONTENTS

INTRODUCTION.....	1
CONTRIBUTING AUTHORS.....	2
PREVIOUS WORK	4
PRODUCT COMPONENTS.....	4
Spatial layers.....	4
Spreadsheets.....	4
MAPPING AND COMPILATION METHODOLOGY.....	4
Field methods.....	4
Compilation methods.....	5
SIGNIFICANT OBSERVATIONS	5
ACKNOWLEDGEMENTS.....	6
REFERENCES	7

LIST OF FIGURES

Figure 1. Location of the Upper Klamath Basin. The red outline shows the extent of the basin in Oregon.....	1
Figure 2. Map of Upper Klamath Basin, Oregon, showing source geologic quadrangle maps.....	3

LIST OF TABLES

Table 1. Geologic quadrangle maps used for Upper Klamath Basin geologic compilation map.....	2
---	---

LIST OF MAP PLATES

Plate 1. Geologic compilation map of part of the Upper Klamath Basin, Klamath County, Oregon	
Plate 2. Fault compilation map of part of the Upper Klamath Basin, Klamath County, Oregon	

INTRODUCTION

The Upper Klamath Basin is located in south-central Oregon (Figure 1). It encompasses the drainage and tributaries of the uppermost watershed of the Klamath River (Figure 2). The river flows out of Oregon into California in the southwest corner of the basin and reaches the Pacific Ocean in northern California. Major tributary rivers to the Klamath within the Upper Basin include the Wood River, the Williamson River, the Sprague River, and the Sycan River. The Lost River also flows through the east side of the basin and in pre-historic times was connected during spring flooding seasons to the Klamath River. It is presently loosely connected to the Klamath River through the irrigation systems of the Upper Klamath Lake Reclamation Project. Several large lakes currently occupying the valley floors are remnants of much larger prehistoric lakes. These lakes include Upper Klamath Lake, Lake Ewana, Round Lake, Aspen Lake, Long Lake, Lower Klamath Lake, Tule Lake, Lake of the Woods, and Swan Lake.

Recent mapping efforts in the basin by the Oregon Department of Geology and Mineral Industries (DOGAMI) began in the late 1990s with new 1:24,000 scale mapping within the Klamath Falls urban growth boundary in support of earthquake hazard analysis. As water issues within the basin became more important, DOGAMI continued making new, large-scale geologic maps in support of the hydrologic studies of the Oregon Department of Water Resources and the U.S. Geological Survey (USGS) Water Resource Department. Over the years seven DOGAMI geologists worked at various times to make new 1:24,000 scale geologic maps. The mapping was principally funded by departmental funding, grants from the Hatfield Committee, funding

from the USGS Statemap program, and various other smaller funding sources. Other recent mapping efforts in the area include the private research in the Cascade Mountains by Stanley Mertzman (Franklin and Marshall College), the USGS sponsored mapping of the Crater Lake National Park by Charles Bacon, and the USGS Edmap program funded mapping and hydrologic studies of the Klamath Marsh area by Michael Cummings and his students at Portland State University.

Work on the present compilation map began in the spring of 2005 with funding from the U.S. Bureau of Reclamation. Supplemental funding was given by the U.S. Forest Service. Reconnaissance field mapping of 17 quadrangles focused on sampling of locatable volcanic eruptive points and sampling and observation of lava flow sections exposed in major fault escarpments. Final reconnaissance-level maps were drawn on the basis of these observations and analysis of aerial photographs. Approximately one week of field time was spent in each quadrangle.




Figure 1. Location of the Upper Klamath Basin. The red outline shows the extent of the basin in Oregon.

CONTRIBUTING AUTHORS

The following lists in alphabetical order the authors who contributed their mapping to this compilation. In parentheses following each quadrangle name is the number of the quadrangle, for example (35), as shown on the location map of the Upper Klamath Basin (Figure 2).

Table 1. Geologic quadrangle maps used for Upper Klamath Basin geologic compilation map.

Contributing Geologist and Affiliation	Quadrangle Map		
	Published	Unpublished	Reconnaissance
Charles Bacon, U.S. Geological Survey		Sun Pass (3) Pothole Butte (1)	
Gerald Black, DOGAMI, retired	Lorella (40) (2004)	Yonna (33) Sprague River West (25) Buttes of the Gods (17)	
Michael Cummings, Portland State University Geology Department		Soloman Butte (9)	
Frank Hladky, formerly with DOGAMI	Bonanza (39) (2003a) Dairy (38) (2003b) Keno (35) (2002)	Agency Lake (14) Chiloquin (15) Beatty (27) Sprague River East (26) Hamaker Mountain (41) Lost River (43)	
Margaret Jenks, DOGAMI	Bryant Mountain (46) (2004) Langell Valley (47) (2004) Malin (45) (2003)	S'ocholis Canyon (16) Worden (42)	Whiteline Reservoir (31) Yainax Butte (34) Swan Lake (32) Swan Lake Point (24) Applegate Butte (10) Cooks Mountain (18) Spodue Mountain (19) Calimus Butte (11) Modoc Point (23) Fuego (4) Lenz (2) Sun Pass (3) Pothole Butte (1) Mares Egg Spring (7) Crystal Spring (13) Pelican Bay (21) Buckhorn Springs (5)
Ian Madin, DOGAMI	Merrill (44) (2003)		
Stanley Mertzman, Franklin and Marshall College Geology Department		Lake of the Woods North (20) Pelican Butte (12) Aspen Lake (28) Devils Peak (6)	
Robert Murray, consulting geologist		Wocus (30)	
George Priest, DOGAMI, retired		Altamont (37) Klamath Falls (36)	
Tom Wiley, DOGAMI	Fort Klamath (8) (2004)	Howard Bay (29) Shoalwater Bay (22)	

DOGAMI is Oregon Department of Geology and Mineral Industries. Two-digit numbers in parentheses are quadrangle numbers in Figure 2. Four-digit numbers are publication years for cited published maps.




Figure 2. Map of Upper Klamath Basin, Oregon, showing source geologic quadrangle maps. Numbers on the figure correspond to geologic quadrangle map numbers in Table 1.

PREVIOUS WORK

Peterson and McIntyre (1970) mapped the eastern part of Klamath County as well as the western part of Lake County at a scale of 1:250,000. They divide the area into 23 units. Their report provides a basic description of the units, a list of fossil localities, and an extensive discussion of the area's mineral resources.

Sherrod and Pickthorn (1992) mapped the west half of the Klamath Falls 1° × 2° quadrangle at a scale of 1:250,000. They divided the area into 30 units and obtained 21 new radiometric rock age dates.

PRODUCT COMPONENTS

In addition to this text, this product includes Geographical Information System (GIS) spatial data layers, map plates created from the GIS data (Plate 1 and Plate 2), and data spreadsheets. File names are in *italic text*.

Spatial layers

- Geology polygon layer (*GEOLOGY*) — Attributed with two different theme labels: map_unit (1:100,000 scale Upper Klamath Basin (UKB) compiled geologic units) and hydro_unit (more generalized lithology/hydrogeology units); see *UKB units legend.xls* spreadsheet, below, for label explanation for each unit
- Fault line layer (*Faults*) — Attributed with quad_name, fault_name (name given to major fault traces), trace_type, side_down (the side of the fault that has moved down relative to the upper fault block), and trace_cert (location of the fault: certain, approximate, or concealed)
- Geochemical samples point layer (*Geochem*) — Attributed with quad_name, sample_no, utm_e and utm_n (location coordinates), iugs_cls (International Union of Geological Scientists [IUGS] volcanic lithology classification type [Le Bas and Streckeisen, 1991]), auth_unit (the contributing author's original geologic unit label), and chem_type (Upper Klamath Basin chemical type; see *UKB chemical types legend.xls* spreadsheet, below, for label explanation for each unit)
- Radiometric age samples point layer (*Age*) — Attributed with sample_no, age_ma (radiometric age date in millions of years), quad_name, source (publication or contributing author's name), and utm_e and utm_n (location coordinates)
- Volcano eruptive point layer (*Volc_loctn*) — Attributed with quad_name, volc_type (general type of volcano: H, hydrovolcanic; V, subaerial vent; or C, cinder cone), volc_name (name given to the topographic feature that is the vent area of the volcano), and utm_e and utm_n (location coordinates)

Spreadsheets

- Legend for geology and lithology/hydrogeology unit polygon labels with unit names and applicable radiometric ages (*UKB units legend.xls*)
- Geochemical analysis results (*UKBgeochemical analyses.xls*)
- Definitions of UKB geochemical volcanic types from the SiO₂, TiO₂, and Al₂O₃ analysis results, their IUGS classification rock type, their comparative relationships (H, high; M, medium; L, low), and their numerical percent ranges (*UKB chemical types legend.xls*)
- Radiometric age date synthesis (*UKB age synthesis.xls*)

MAPPING AND COMPILATION METHODOLOGY

Field methods

Contributing authors used a variety of methods for their field work. These methods include walking out contacts between rock units of different types, sampling and making observations at specific locations, locating samples with GPS and field computers as well as on paper topographic maps, and writing logs of observations and sample locations. Field observations are enhanced by observation of aerial photographs, analyses of sample rocks for their geochemical compositions and their radiometric ages, and thin section study of rock samples. The original geologic maps, before compilation, were all completed on 1:24,000 scale USGS standard topographic base maps.

Compilation methods

The only equivalent data available throughout the study area were geochemical analyses. To compare analyses, first, all analyses were normalized and plotted on the IUGS volcanic rock classification (Le Bas and Streck-eisen, 1991) diagram ($\text{SiO}_2/\text{Total alkali}$). Several different elemental comparisons were then tried, including Ba/SiO_2 , Sr/SiO_2 , Ba/K , and Sr/K . None yielded consistent lithology group results throughout the database. Finally, analyses were sorted by K_2O within each IUGS volcanic rock classification. The amounts of SiO_2 , TiO_2 , and Al_2O_3 were compared and classified according to their relative percentages (high, medium, low). This methodology created generalized classes of volcanic rocks, which were assigned alphanumeric labels (for example, Tand6 and Tbas10). These classes were then plotted spatially and compared to the original author's map units. A unit label, composed of the age of the unit from the radiometric ages and stratigraphic relations in the vicinity and the informal geologic unit name, was assigned for each area of similar unit and class type. Lithologic/hydrogeologic units were assigned only on the basis of age and IUGS rock classification type.

SIGNIFICANT OBSERVATIONS

The compiler made the following list of observations on the basis of her published, unpublished, and reconnaissance mapping and her compilation of other authors' observations and mapping. Although most of the observations resulted from the compiler's direct field observations, they are not the conclusions or opinions of the other authors who contributed geologic mapping and information to the project.

1. Volcanic activity within the basin has been nearly continuous during the last 8 million years, with no significant breaks, given our present understanding of radiometric ages of the basin's volcanic rocks.
2. A major part of the area is underlain by volcanic rocks and sediments ranging in age from 2.0 to 5.0 Ma (Pliocene).
3. The seemingly pervasive sequences of subaerial lava flows underlying the topographic surfaces throughout the basin are recent, thin caps of lava flows over diverse sections of thick older sediments, hydrovolcanic deposits, and water-affected lava flows.
4. Faults, especially large valley bounding faults, are made up of a number of parallel fault planes.
5. The parallel fault planes successively downdrop the capping volcanic units, creating repeated sections in the fault escarpments.
6. Faulting not only creates steep escarpments but also tilts the upper surfaces of the lava flows or sediments up and away from the fault escarpment.
7. Faulting of the basin by normal faults is fairly recent, beginning 1.5 to 2.0 million years ago.
8. Fault escarpments show very little stream dissection, and the valley floor alluvial fans at the bases of the escarpments are small.
9. Previous to the late Pliocene/early Pleistocene faulting the landscape was fairly level with some low hills of older calc-alkaline volcanic rocks surrounded by the flows of the younger basaltic rocks.
10. Most of the basin's volcanic flows and sedimentary and volcanic detritus principally spread out over nearly horizontal ground surfaces.
11. Valley filling by lava flows is found only in the northern part of the area, particularly in relation to the Sprague River and Sycan River ancient drainages.
12. Valley-confined lava flows filled broad, shallow valleys rather than steep-sided, deep canyons.
13. Faulting in the southern part of the basin is more intense and extensive, with dominant structural grains including northwest-southeast, east-west, and north-south.
14. With the exception of the northern part of the Upper Klamath Lake graben, faulting in the northern part of the basin is less intense and extensive, with smaller displacements on the fault escarpments and a general north-south structural grain.
15. Volcanic lithologies are variable, ranging from rhyolites to trachybasalts.
16. Volcanic sequences, trending upward from calc alkaline to more primitive tholeiitic, are repeated across the area from east to west and south to north.
17. The volcanic activity becomes younger from east to west.

18. The volcanic sequences are repeated through time from east to west.
19. Nearly 470 volcanic vents are located in the basin area.
20. In volume, the eruptive products from the individual volcanic vents are not large and are localized around the vent/vents that erupted them.
21. With the exception of Mount Mazama, the volcanoes in the area were erupted as single-episode events, confined to specific, short periods (tens of thousands to hundreds of thousands of years)
22. Stream and river drainage locations are determined by faulting or lava flow edges.
23. Significant deposition of finer grained, lake and low-gradient stream sediments has occurred throughout the basin's geologic history.
24. Interlayered with the sediments throughout the geologic history of the basin are hydrovolcanic units including hydroclastic cinder cones, water affected basalt flows, and pillowed basalt flows.
25. Significant erosion occurred following the deposition of sediments and hydrovolcanic units and before the deposition of the capping subaerial lava flows.
26. Much of the present topography on the south side of the Sprague River Valley is exhumed terrain.
27. Most of the hydrovolcanic eruption exposures have been removed by erosion, leaving only the volcano's cemented central vent "root" deposits.
28. Glaciation in the Cascade Range on the west side of the area was extensive and sheetlike.
29. Glaciation in the Cascade Range was disrupted by both faulting and subsequent post-glacial valley-formation lava flows.
30. Faulting on the west side of the Upper Klamath Lake is younger than the end of the latest glacial period and cuts moraine and alluvial fan deposits in numerous locations.
31. Above the elevation of approximately 1,800 m (6,000 ft), valley glaciers have carved the landscape throughout the area east of Upper Klamath Lake.
32. Valley glaciers have followed fault valleys and escarpments as well as lava flow edges.
33. Most larger alluvial fans in the basin are actually glacial fans formed by glacial debris at the ends of the valley glaciers.
34. Most springs, both hot and cold water, in the basin are located along valley bounding faults.
35. The Mazama ash deposits on the east side of the Klamath Marsh blanket but do not fill a pre-existing terrain of lava flows and dissected sediments.
36. Subsurface water flows within the Mazama ash deposits are directed in part by the underlying pre-existing terrain.
37. Pre-existing glacially eroded valleys on the east side of the Mazama volcano were the pathways/guideways for the Mazama eruption ash flows.
38. Large pothole features, found in the Mazama deposits in three separate areas, may be the result of the post-eruption melting of large blocks of glacial ice.
39. A well-established, post-eruption, low-gradient drainage pattern on the west side of the Klamath Marsh has been affected by post-eruption faulting and has been replaced with a present-day incisional drainage pattern.
40. Fish fossils from the mapped area all yield relative ages of 4.0 to 6.0 Ma (Pliocene), with a juvenile spawning population found in one locality.

ACKNOWLEDGEMENTS

This study was partially funded by U.S. Bureau of Reclamation contract number 04 PG250049, titled Klamath Basin Geological Compilation, and U.S. Department of Agriculture, U.S. Forest Service, project AG-04H1-P-05-0005.

The compiler thanks the Winema National Forest soils group for loaning aerial photographs for photocopying; Lina Ma, for field assistance and geochemical sample normalization analysis; Marina Drazba, for digitizing of reconnaissance mapping; Paul Staub, for GIS quadrangle appending, geometry management, and metadata; John Huard, Oregon State University, for analysis of flow age samples; Paul Layers, University of Alaska, Fairbanks, for analysis of ash and glass age samples; Stan Mertzman, Franklin and Marshall University, for geochemical sample analysis; Gerald Smith, University of Michigan, Ann Arbor, for fish fossil analysis; Michael Cummings, Portland State University, for geologic discussions; Jeld-Wen Corporation forestry department, for property access; private land-owners of Klamath County, for property access and local history; Charles Cossey, for field, transportation, and lodging assistance; and David Sherrod, USGS, for field maps for the 1992 Klamath Basin geologic map.

REFERENCES

- Black, G., 2004, Geologic map of the Lorella quadrangle, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Open-file Report O-04-07, CD, scale 1:24,000.
- Hladky, F. R., and Mertzman, S., 2002, Geologic map of the Keno quadrangle, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Geological Map Series GMS-102, 17 p., CD, scale 1:24,000.
- Hladky, F. R., 2003a, Geologic map of the Bonanza quadrangle, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Geological Map Series GMS-116, 22 p., CD, scale 1:24,000.
- Hladky, F. R., 2003b, Geologic map of the Dairy quadrangle, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Geological Map Series GMS-112, 20 p., CD, scale 1:24,000.
- Jenks, M. D., 2004, Geologic map of the Bryant Mountain and Langell Valley quadrangles, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Geological Map Series GMS-117, 36 p., CD, scale 1:24,000.
- Jenks, M. D., and Madin, I. P., 2003, Draft geologic map of the Merrill and Malin quadrangles, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Open-file Report O-03-03, CD, scale 1:24,000.
- Le Bas, M. J., and Streckeisen, A. L., 1991, The IUGS systematics of igneous rocks: London, *Journal of the Geological Society*, v. 148, p. 825–833.
- Malin, K., and Hart, W. K., 1991, New K-Ar ages of mafic lavas from the Basin and Range–Cascade transition zone in northeastern California and southern Oregon: *Isochron West*, no. 57, p. 8–11.
- McKee, E. H., Duffield, W. A., and Stern, R. J., 1983, Late Miocene and early Pliocene basaltic rocks and their implications for crustal structure, northeastern California and south-central, Oregon: *Geological Society of America Bulletin*, v. 94, no. 2., p. 292–304.
- Peterson, N. V., and McIntyre, J. R., 1970, The reconnaissance geology and mineral resources of eastern Klamath County and western Lake County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Bulletin 66, 70 p., 2 plates, scale 1:250,000.
- Sherrod, D. R., and Pickthorn, L. B. G., 1992, Geologic map of the west half of the Klamath Falls 1° × 2° quadrangle, south-central Oregon, Reston, VA, US Geological Survey Miscellaneous Investigations Series Map I-2182, 1 sheet, scale 1:250,000.
- Wiley, T. J., 2004, Geologic map of the Fort Klamath quadrangle, Klamath County, Oregon, Portland, OR: Oregon Dept. of Geology and Mineral Industries Geological Map Series GMS-96, 24 p., CD, scale 1:24,000.