

OREGON GEOLOGY

published by the

Oregon Department of Geology and Mineral Industries



VOLUME 43, NUMBER 8

AUGUST 1981



OREGON GEOLOGY

(ISSN 0164-3304)

VOLUME 43, NUMBER 8

AUGUST 1981

Published monthly by the State of Oregon Department of Geology and Mineral Industries (Volumes 1 through 40 were entitled *The Ore Bin*).

Governing Board

John L. Schwabe, Chairman Portland
C. Stanley Rasmussen Baker
Allen P. Stinchfield North Bend

State Geologist Donald A. Hull

Deputy State Geologist John D. Beaulieu

Editor Beverly F. Vogt

Main Office: 1005 State Office Building, Portland 97201, phone (503) 229-5580.

Baker Field Office: 2033 First Street, Baker 97814, phone (503) 523-3133.

Howard C. Brooks, Resident Geologist

Grants Pass Field Office: 312 S.E. "H" Street, Grants Pass 97526, phone (503) 476-2496.

Len Ramp, Resident Geologist

Mined Land Reclamation Program: 1129 S.E. Santiam Road, Albany 97321, phone (503) 967-2039.

Paul F. Lawson, Supervisor

Subscription rates: 1 year, \$4.00; 3 years, \$10.00. Single issues, \$.40 at counter, \$.75 mailed.
Available back issues of *The Ore Bin*: \$.25 at counter, \$.50 mailed.

Address subscription orders, renewals, and changes of address to *Oregon Geology*, 1005 State Office Building, Portland, OR 97201.

Send news, notices, meeting announcements, articles for publication, and editorial correspondence to the editor, Portland office. The Department encourages author-initiated peer review for technical articles prior to submission. Any review should be noted in the acknowledgments.

Permission is granted to reprint information contained herein. Credit given to the Oregon Department of Geology and Mineral Industries for compiling this information will be appreciated.

Second class postage paid at Portland, Oregon.

Postmaster: Send address changes to *Oregon Geology*, 1005 State Office Building, Portland, OR 97201.

COVER PHOTO

Mafic dike of John Day Formation cutting Cretaceous marine mudstones of the Hudspeth Formation along Gable Creek in the Mitchell quadrangle, Oregon. The article beginning on the next page discusses these dikes.

New index of Oregon topo maps available from USGS

A new index showing areas of the State of Oregon covered by 1,091 published topographic maps is now available upon request from the U.S. Geological Survey (USGS).

By showing the shape and elevation of the terrain and delineating a wide range of natural and manmade features, standard USGS topographic quadrangle maps are valuable records of the land surface for engineers, scientists, planners, and others concerned with the nation's resources. The maps also are popular with hikers, hunters, campers, and other open-air enthusiasts as "silent guides" to the outdoors. The USGS annually distributes more than 10.5 million copies of its 45,000 published "topo" maps.

The Oregon index includes a map of the state showing areas covered by each of the available 7½-minute and 15-minute quadrangle maps. The text on the back of the index lists other maps of Oregon published by the Survey, such as maps of the entire state, rivers and creeks, reservoirs and dam-sites, and Crater Lake National Park.

More than 85 percent of Oregon has now been covered by quadrangle maps. The 7½-minute maps are at a scale of 1:24,000 (1 in. equals 2,000 ft) and the 15-minute quadrangle maps are at a scale of 1:62,500 (1 in. equals about 1 mi).

USGS-published maps of Oregon are deposited and available for inspection at nine reference libraries in the state—in Ashland, Bend, Corvallis, Eugene, Klamath Falls, La Grande, Monmouth, Portland, and Salem. The index also lists commercial outlets in Oregon, California, Idaho and Washington where many of the maps may be purchased, including the Portland office of the Oregon Department of Geology and Mineral Industries.

Copies of the "Index to Topographic Maps of Oregon" are available free from USGS Public Inquiries Offices in Menlo Park, Calif. (Building 3, 345 Middlefield Rd.); San Francisco (504 Custom House, 555 Battery St.); Los Angeles (7638 Federal Building, 300 North Los Angeles St.) and Spokane, Wash. (678 U.S. Court House, West 920 Riverside Ave.). □

Mine reclamation fees increase

The 1981 Oregon Legislative Assembly passed House Bill 2220 increasing the fees (provided in ORS 517.800) for mining reclamation. The bill was signed by the Governor and became effective upon signature July 2, 1981.

The application fee for a new site is \$390, and the annual renewal fee is \$290. The increase is designed to maintain the self-sufficiency of the Mined Land Reclamation Program through the 1981-1983 biennium. This program, intended to restore surface mined land to a continuing useful purpose, is 90 percent funded by the industry it regulates.

The new fees are now effective for all new mining site applications. Renewal fees are effective August 1981. □

CONTENTS

A mafic dike system in the vicinity of Mitchell, Oregon, and its bearing on the timing of Clarno-John Day volcanism and early Oligocene deformation in central Oregon	107
Oregon oil and gas activity for first half of year	113
Abstract	114
Engineering Geologist meeting set for Portland	114

A mafic dike system in the vicinity of Mitchell, Oregon, and its bearing on the timing of Clarno-John Day volcanism and early Oligocene deformation in central Oregon

by Edward M. Taylor, Department of Geology, Oregon State University, Corvallis, Oregon 97331

INTRODUCTION

A system of west-northwest-trending basaltic dikes crops out in the vicinity of Mitchell, Oregon. The dikes are exposed best in the central part of the northeast-southwest-trending Mitchell anticline, where they have intruded Cretaceous marine sedimentary rocks of the Gable Creek and Hudspeth Formations. The dikes also have penetrated the overlying volcanic rocks of the Clarno Formation on both limbs of the anticline and were probably associated with basaltic lava flows of the lower John Day Formation. Parts of this dike system appear on a published geologic map of the Mitchell quadrangle and have been designated "Airport dikes," "Nelson Creek dikes," and "Keyes Creek dikes" (Oles and Enlows, 1971).

The mafic dike system has been traced over an area ap-

proximately 9 mi long and 2 mi wide in which the dikes are arranged in groups of subparallel segments (Figure 1). Individual segments range in length from 1 mi to a few tens of feet and are commonly offset from each other along trend. Adjacent segments locally taper to thin edges at their extremities, or they are joined, nearly at right angles, by short connective dikes. Most of the dike segments are 15-20 ft wide; their width at a few localities where bulbous protrusions extend into incompetent rocks is as much as 110 ft. In contrast, some dikes apparently were unable to invade competent rock; they terminate or bifurcate just below thick resistant strata or preexistent sills.

Elongate depressions are produced by erosion of dikes that cut resistant strata (Figure 2). Elsewhere, as in soft mudstones of the Hudspeth Formation, low ridges mark the positions of dikes (Figure 3). Volcanic mudflow deposits of the

Figure 1. Distribution of high Fe-Ti tholeiitic basalt dikes of late Oligocene age in the vicinity of Mitchell, Oregon. Heavy solid lines represent dike outcrops; dotted lines represent inferred positions of covered dikes.






Figure 2. Mafic dike in negative relief. Country rock is erosionally resistant conglomerate of the Gable Creek Formation, 3.8 mi west of Mitchell.

Clarno Formation have been so indurated in close proximity to the dikes that double walls of erosionally resistant mudflow can be found standing in relief along dike margins.

The margins of dikes consist of very fine-grained, black, devitrified glass in a chilled zone up to 2 ft wide. Dike interiors are gray where fresh, uniformly coarse grained, and commonly display crude columnar jointing (Figure 4). Weathering of dike rocks is always more advanced in the interior than at the margins and produces a dark-brown, ferruginous soil.

PETROGRAPHY

In the dike margins, thin tablets of plagioclase (An_{55}) make up 3-4 percent of the rock and are never more than 0.5 mm long. They are surrounded by grains of pyroxene and magnetite approximately 0.001 mm in diameter which form a very fine-grained intergranular texture with microlites of plagioclase. Thin films of brown glass still exist in the freshest samples. A green isotropic clay mineral, probably a complex chlorite interlayered with other phyllosilicates replacing traces of olivine, occurs in euhedral patches up to 0.5 mm in diameter (Figure 5A).

The zone of transition between the fine-grained margins and the coarse-grained interior of dikes is only 7-10 in. wide (Figure 6). Within this zone, elongate crystals of clinopyroxene and plagioclase are suspended in a fine groundmass and become larger and more abundant toward the interior. Near the inner boundary of the transition zone, the groundmass is represented by small, fine-grained patches of late-crystallizing



Figure 3. Mafic dike in positive relief. Country rock is easily eroded mudstone of the Hudspeth Formation, 2.8 mi west of Mitchell.



Figure 4. Columnar jointing in the Keyes Creek dike adjacent to U.S. Highway 26. Hat rests upon contact with volcanic mudflow deposits of the Clarno Formation.

feldspar, largely altered to smectite (Figure 5B).

Wherever the width of dikes exceeds 10-12 ft, the interior is texturally hypidiomorphic-granular, consisting of plagioclase, monoclinic pyroxene, and opaque oxides up to 1.5 mm in length (Figure 5C). Plagioclase of composition An_{50-60} makes up 75 percent of the dike rocks. The pyroxene is a dusky augite with a 2V of 45-48° and commonly constitutes 18-20 percent of the rock. The opaque oxide, which is titanomagnetite in the fine-grained margin, becomes a coarse-grained skeletal and dendritic admixture of ilmenite and magnetite in the interior.

Alteration of dike rocks by deuteric processes and weathering has converted all olivine to green chloritic minerals. In varying degrees, pyroxene is altered to highly birefringent yellow and orange smectite, ilmenite and magnetite are altered to leucoxene and hematite, and plagioclase is replaced by carbonate. In Table 1, the chemical effects of alteration are revealed by the contrast between column 3, an average of two very fresh samples from dike margins, and column 4, an average of seven altered samples from dike interiors. Altered dike rocks have lost Fe and Mg but gained Ca and an unspecified quantity of H_2O and CO_2 .

Rocks of the mafic dike system in the Mitchell area are low-alumina tholeiites enriched in Fe and Ti. They are distinct from Mitchell-area basaltic flows and dikes of the Clarno Formation in which the Al, Ca, and Mg content is much greater and the Fe and Ti content is much smaller (Taylor, unpublished data). Major-element composition of the dikes is also distinct from basalts of the Picture Gorge Formation in the Mitchell area because the dikes contain much more Fe and Ti. The closest compositional match to the mafic dike system is

Table 1. Major-element composition of mafic dikes near Mitchell, Oregon*

	(1)	(2)	(3)	(4)	(5)
SiO ₂	51.3	51.8	51.6	52.0	47.74
Al ₂ O ₃	12.8	13.4	13.1	13.7	15.27
FeO	15.6	14.5	15.1	14.1	15.2
CaO	7.8	8.5	8.2	8.6	7.57
MgO	5.4	4.0	4.7	3.5	4.70
K ₂ O	0.65	0.50	0.60	0.75	1.46
Na ₂ O	3.1	3.1	3.1	3.0	3.48
TiO ₂	3.10	3.00	3.05	3.00	3.33
Total	99.75	98.80	99.45	98.65	98.75

- (1) Fresh marginal glass from Keyes Creek dike, 500 ft north of south corner secs. 31 and 32, T. 11 S., R. 22 E.
- (2) Fresh marginal glass from east end of Mill Creek dike, 4,400 ft elevation, NW ¼ sec. 9, T. 12 S., R. 22 E.
- (3) Average of (1) and (2).
- (4) Average of seven slightly altered rocks from dike interiors, representing all dike groups.
- (5) Basaltic lava from lower John Day Formation, 6 mi northwest of Mitchell, 700 ft west of intersection of secs. 4, 5, 8, and 9, T. 11 S., R. 21 E. Recast H_2O -free from Hay (1962).

* Analyses (1)-(4) by XRF and AAS. Total Fe as FeO. H_2O and CO_2 not included because samples were fused before analysis.




Figure 5. Photomicrographs of dike rocks in which each field width is 0.2 mm. (A) Fine-grained margin consisting of sparse olivine altered to clay (center) surrounded by microlites of feldspar and granules of pyroxene, titanomagnetite, and glass. (B) Transition zone of coarse feldspar, monoclinic pyroxene, blades of intergrown magnetite and ilmenite, and patches of fine groundmass. Crossed polars. (C) Coarse hypidimorphic-granular interior. Same minerals as (B).



Figure 6. Transition zone from fine-grained, close-fractured margin (below head of hammer) to coarse-grained interior of mafic dike (above head of hammer).

with basaltic lavas in the lower part of the John Day Formation northwest of Mitchell (Figure 1). The John Day lavas lack phenocrysts and contain high Fe and Ti (Table 1, column 5). It is not suggested that the mafic dike system was the direct source of these particular John Day lavas, because the dikes are high-Si, low-K tholeiites while the lavas are low in Si and high in K, transitional to alkali basalts (Robinson, 1969).

CONCLUSIONS AND SPECULATIONS

Emplacement of the dikes

An obvious feature in need of resolution is the rather peculiar distribution of the dikes. Within the overall west-northwest trend, the dikes occur in distinct groups that are successively offset en echelon in a left-lateral sense. For example, Nelson Creek dikes seem to be offset from Mill Creek dikes, and a similar pattern is exhibited by Gable Creek, The Narrows, Airport, and Keyes Creek dikes (Figure 1). On a smaller scale, the same pattern can be seen between the individual off-set segments of each dike group.

Although it has been reported that "dikes generally occupy preexisting faults" and some dike segments have been shown on geologic maps to be offset by strike-slip faults (Oles and Enlows, 1971, map and explanation), detailed study of the dikes and surrounding country rock has failed to confirm these

relationships. Many faults of varying extent and displacement occur in the vicinity of dikes but not one has been found to be coincident with a dike. Slickensided fractures have been found within dike rocks at two localities; displacement of only a few feet is indicated. Where extremities of dikes are exposed, they are seen to taper to thin edges rather than blunt, faulted terminations. Moreover, adjacent segments often overlap in parallel alignment far beyond the positions of possible cross faults.

The Keyes Creek and Nelson Creek dikes have been described as parts of a single dike that has been offset 22,000 ft by right-lateral slip along the Mitchell fault (Oles and Enlows, 1971, p. 48, 54, 59). This interpretation must be reconsidered, because the eastern 1 mi of the Keyes Creek dike cuts across the Mitchell fault (Figure 1). Segments of the dike lie in the crush zone of the fault and display no evidence of disruption or reorientation. If the Nelson Creek dikes are correctly projected to the Mitchell fault, an offset of 28,000 ft is indicated. This is more than twice the right-lateral displacement of the Mitchell fault, as demonstrated by offset of the axis of the Mitchell anticline (Wilkinson and Oles, 1968). It is my conclusion that the mafic dike system in the Mitchell area has not been significantly displaced by later faulting of any kind. In particular, right-lateral movement along the Mitchell fault preceded dike emplacement.

The dike segments are not randomly located; in both position and orientation their emplacement appears to have been systematically controlled. It is especially significant that the dikes, with one exception, seem to have avoided the trace of the Mitchell fault and many other faults that are parallel to it. What could have localized the dikes? This question can be readily answered if it is assumed that the dikes were emplaced after most of the right-lateral movement had occurred on the Mitchell fault but before the responsible conditions of stress had died out. The magma would have found paths of least resistance along gash fractures at acute angles to the Mitchell fault, while passage of magma into the fault zone itself would have been hindered. Gash fracture control of the dikes was proposed by Oles and Enlows (1971, p. 59), but they also em-

phasized dike emplacement prior to right-lateral faulting and presumably prior to development of gash fractures.

A new emphasis should be placed upon the nearly equivalent age of the dike system and the Mitchell fault. If a deep-seated body of mafic, nonporphyritic, fluid magma, elongate in a west-northwest direction, was intersected by a zone of right-lateral disruption of the crust along the Mitchell fault, the magma could have penetrated gash fractures and produced the pattern of dikes now seen in the Mitchell area. Indeed, without this combination of tectonic pressures and avenues of escape, the dense magma might never have reached shallow crustal levels or poured over the surface.

Age of the dikes

K-Ar ages of the Nelson Creek dikes and Airport dikes were reported to be 29.4 ± 0.6 and 33.4 ± 1.3 m.y., respectively (Enlows and Parker, 1972). However, the dikes were probably formed at the same time. It is unlikely that, in this small area, many separate magmas of nearly identical major-element composition and extent of crystallization would penetrate to the same crustal level in response to apparently identical conditions of stress and adopt the same paleomagnetic polarity over a span of four million years. Because the dikes show a much closer compositional relationship to lavas of the John Day Formation than to any known rocks of the Clarno Formation, and because the age (31.5 m.y.) of John Day mafic lavas only 3 mi northwest of the dikes (Evernden and others, 1964) matches the average age (31.4 m.y.) of dikes reported by Enlows and Parker, I conclude that 31.5 m.y. is the best currently available estimate of the age of the mafic dike system in the Mitchell area.

Clarno-John Day boundary problem

Enlows and Parker (1972) placed the dikes within the "indicated span of Clarno igneous activity in the Mitchell quadrangle" which lasted "from about 46 million years before pres-

Table 2. K-Ar ages from upper Clarno and lower John Day Formations in the vicinity of Mitchell, Painted Hills, and Clarno Bridge

Age (m.y.)	Material	Location	Stratigraphic position	Reference
29.3 ± 0.5 29.4 ± 0.6	Whole rock	1 mi SE of Mitchell	Nelson Creek dikes	Enlows and Parker, 1972
31.1	Sanidine	Painted Hills	Tuff bed 165 ft above base of John Day Formation	Evernden and others, 1964
31.5	Whole rock	Between Mitchell and Painted Hills	Basalt lava in lower red John Day Formation	Evernden and others, 1964
32.0	Sanidine	1 mi E of Clarno Bridge	Altered tuff at base of John Day Formation	Evernden and James, 1964
30.1 ± 4.7 35.6 ± 3.4	Whole rock	4.5 mi E of Mitchell	Andesite lava, "upper Clarno"	Enlows and Parker, 1972
33.3 ± 1.2 33.5 ± 1.3	Whole rock	1.5 mi NW of Mitchell	Airport dikes	Enlows and Parker, 1972
34.0	Plagioclase	2 mi NE of Clarno Bridge	"Nut Beds" at top of Clarno Formation	Evernden and James, 1964
36.4 ± 1.1	Sanidine	2.2 mi SW of Ashwood	"Member A" at base of John Day Formation	Swanson and Robinson, 1968
36.5 ± 0.9	Sanidine	1.5 mi SW of Painted Hills	Crystal tuff near top of Clarno Formation	Evernden and others, 1964
37.5	Whole rock	Between Mitchell and Painted Hills	Andesite lava near top of Clarno Formation	Evernden and others, 1964

ent to 30 million years before present" and "was immediately succeeded by the deposition of the extensive volcanoclastic sediments of the John Day Formation" (p. 105 and Table 1). This interpretation was supported by an average age of 32.8 m.y. for a Clarno lava sample (KFO-901) collected 4.5 mi east of Mitchell.* In opposition to this interpretation was the 36-m.y. age of a widespread welded ash-flow sheet in "member A" at the base of the John Day Formation (Swanson and Robinson, 1968). Table 2 lists published K-Ar ages of Clarno and John Day rocks pertinent to this boundary problem. Much discussion in recent years has centered upon the possibility of simultaneous Clarno and John Day volcanism in central Oregon.

Inconsistencies between published radiometric ages and known stratigraphic relationships can be removed by reconsideration of three K-Ar determinations. "Member A" (36.4 m.y.) at the base of the John Day Formation overlies bedded John Day tuff (32.0 m.y.) and bedded Clarno tuff (34.0 m.y.) at Clarno Bridge (Evernden and James, 1964; Evernden and others, 1964). These bedded tuffs have been altered to bentonitic clay, and, as suggested by Swanson and Robinson (1968), their feldspars could have lost argon, giving rise to younger ages. The basal ash-flow tuff of "member A," in contrast, is a relatively fresh unit, and its age is probably more reliably estimated. The age of the "upper Clarno" andesite lava (KFO-901) from Mitchell (av. 32.8 m.y.) is erroneous because, in my opinion, it belongs stratigraphically in the lower Clarno Formation. An unpublished K-Ar age of 48.9 ± 5.2 m.y. (Enlows, Robinson, and McKee, personal communication) has been obtained on hornblende separated from another andesite flow in a nearly equivalent stratigraphic position, 3.4 mi northeast of Mitchell. In addition, volcanic mudflow deposits at approximately the same stratigraphic level as the nearby 48.9-m.y. hornblende andesite lava have been intruded by the mafic rocks of Marshall Butte. The average of two Marshall Butte age measurements is 44.9 m.y. (Enlows and Parker, 1972). It is my conclusion that in the vicinity of Mitchell, Painted Hills, and Clarno Bridge, deposition of the Clarno Formation ceased and was rapidly followed by deposition of the John Day Formation approximately 36 million years ago.

Early Oligocene deformation

The lower ash-flow tuff of "member A" at the base of the John Day Formation appears to have spread across the trace of the Blue Mountain anticline eastward to within 13 mi of Mitchell (Robinson, 1975). Most of the later John Day ash-flow tuffs were prevented from spreading that far to the east by a topographic barrier coincident with the Blue Mountain anticline (Robinson, 1966). Uplift of the Blue Mountain anticline probably produced the topographic barrier after much of the lower John Day Formation had been deposited. These relationships are compatible with an episode of northwest-southeast compressional deformation of sufficient intensity to produce broad folds and strike-slip faults and to release relatively small volumes of deep-seated mafic magmas to the surface during early Oligocene time in north-central Oregon.

The obvious magmatic distinctions between Clarno and John Day volcanism developed independently from, and some five million years prior to, the onset of crustal deformation that produced folds in John Day rocks. It is suggested that lower John Day tuffs will be found resting conformably on

upper Clarno rocks in some localities and that an angular unconformity will be recognized between lowermost John Day tuffs and overlying John Day tuffs where preservation and exposure permit careful measurement. In similar fashion, strike-slip movement on the Mitchell fault probably displaced lower John Day tuffs but not overlying members of the John Day Formation.

ACKNOWLEDGEMENTS

My appreciation is extended to K.F. Oles and H.E. Enlows for their assistance in reading the manuscript of this paper.

REFERENCES CITED

- Enlows, H.E., and Parker, D.J., 1972, Geochronology of the Clarno igneous activity in the Mitchell quadrangle, Wheeler County, Oregon: Oregon Department of Geology and Mineral Industries, Ore Bin, v. 34, no. 6, p. 104-110.
- Evernden, J.F., and James, G.T., 1964, Potassium-argon dates and the Tertiary floras of North America: American Journal of Science, v. 262, no. 8, p. 945-974.
- Evernden, J.F., Savage, D.E., Curtis, G.H., and James, G.T., 1964, Potassium-argon dates and the Cenozoic mammalian chronology of North America: American Journal of Science, v. 262, no. 2, p. 145-198.
- Hay, R.L., 1962, Origin and diagenetic alteration of the lower part of the John Day Formation near Mitchell, Oregon, in *Petrologic studies—A volume in honor of A.F. Buddington*: New York, N.Y., Geological Society of America, p. 191-216.
- Oles, K.F., and Enlows, H.E., 1971, Bedrock geology of the Mitchell quadrangle, Wheeler County, Oregon: Oregon Department of Geology and Mineral Industries Bulletin 72, 62 p.
- Robinson, P.T., 1966, Facies changes in John Day Formation, Oregon (abs.): Geological Society of America Abstracts for 1966, GSA Special Paper 101, p. 417.
- — — 1969, High-titanium alkali-olivine basalts of north-central Oregon, U.S.A.: Contributions to Mineralogy and Petrology, v. 22, no. 4, p. 349-360.
- — — 1975, Reconnaissance geologic map of the John Day Formation in the southwestern part of the Blue Mountains and adjacent areas, north-central Oregon: U.S. Geological Survey Miscellaneous Investigations Series Map I-872.
- Swanson, D.A., and Robinson, P.T., 1968, Base of the John Day Formation in and near the Horse Heaven mining district, north-central Oregon: U.S. Geological Survey Professional Paper 600-D, p. D154-D161.
- Wilkinson, W.D., and Oles, K.F., 1968, Stratigraphy and paleoenvironments of Cretaceous rocks, Mitchell quadrangle, Oregon: American Association of Petroleum Geologists Bulletin, v. 52, no. 1, p. 129-161. □

Check your address label

During the next few months, we will be using a new addressing system to send you your copy of *Oregon Geology*. If you see any errors in your address or encounter any problems in receiving your magazine, please notify us. □

* It should be recorded that the location of KFO-901 is given incorrectly by Enlows and Parker; it was taken from the summit of hill 4703, in the SW ¼ of sec. 35, T. 11 S., R. 21 E. (K.F. Oles, personal communication).

Oregon oil and gas activity for first half of year

The 1980 oil and gas exploration and drilling activity in Oregon was summarized earlier this year (*Oregon Geology*, March 1981). Since the first of the year, however, 13 permits to drill have been granted, with an additional four pending at this time. Seven wells have been drilled this year, for a total of 34,589 ft.

Demand for drilling rigs in other parts of the country has held Oregon's rig count to one or two during the spring months, but an additional rig is expected for the remainder of

the year. Drilling so far in 1981 has been confined to the western part of the state, with the exception of one hole in Crook County in central Oregon. Operators have included American Quasar Petroleum Company; Diamond Shamrock Corporation; John Miller; and Texaco, Incorporated.

In March, American Quasar discovered Oregon's second gas field a mile west of Lebanon in the Willamette Valley. The well, Hickey 9-12, produces at a rate of 200,000 cubic feet per day from sands at a depth of 3,000 ft.

Oil and gas permits and drilling activity since January 1981

Permit number	Operator	Well name	Location	PTD: Proposed (ft) TD: Total depth (ft)	Status
176	American Quasar Petroleum Co.	Franbea 36-34	SE¼ sec. 36 T. 7 N., R. 5 W. Columbia Co.	PTD 5,000	Permitted 1/21/81
177	Diamond Shamrock Corporation	Boise Cascade 11-14	NW¼ sec. 14 T. 5 N., R. 7 W. Clatsop Co.	TD 7,864	Abandoned dry hole
178	Diamond Shamrock Corporation	Crown Zellerbach 11-28	NW¼ sec. 28 T. 5 N., R. 9 W. Clatsop Co.	TD 5,700	Abandoned dry hole
179	Diamond Shamrock Corporation	Crown Zellerbach 31-17	NE¼ sec. 17 T. 6 N., R. 8 W. Clatsop Co.	PTD 10,000	Drilling
180	Reichhold Energy Corporation	Columbia Co. 32-10	NE¼ sec. 10 T. 6 N., R. 5 W. Columbia Co.	PTD 8,000	Permitted 3/3/81
181	Reichhold Energy Corporation	Columbia Co. 23-5	SW¼ sec. 5 T. 6 N., R. 5 W. Columbia Co.	PTD 3,500	Permitted 3/12/81
182	Reichhold Energy Corporation	Columbia Co. 13-1	SW¼ sec. 1 T. 6 N., R. 5 W. Columbia Co.	PTD 3,500	Permitted 3/12/81
183	Reichhold Energy Corporation	Hemeon 14-14	SW¼ sec. 14 T. 6 N., R. 5 W. Columbia Co.	PTD 2,700	Permitted 3/12/81
184 ^{ac}	Reichhold Energy Corporation	Longview Fibre 41-32	NE¼ sec. 32 T. 7 N., R. 5 W. Columbia Co.	PTD 3,200	Permitted 4/10/81
185	American Quasar Petroleum Co.	Kenneth Wetgen 26-32	NE¼ sec. 26 T. 13 S., R. 4 W. Linn Co.	PTD 5,000	Permitted 4/2/81
186	American Quasar Petroleum Co.	Wolverton 13-31	NE¼ sec. 13 T. 10 S., R. 3 W. Marion Co.	PTD 5,000	Permitted 4/21/81
187	Reichhold Energy Corporation	Ellis 23-26	SW¼ sec. 26 T. 5 N., R. 4 W. Linn Co.	PTD 4,000	Permitted 4/29/81
188	American Quasar Petroleum Co.	Chipman 4-14	SW¼ sec. 4 T. 12 S., R. 2 W. Linn Co.	PTD 5,000	Permitted 6/09/81
	Miller Drilling Co.	Bork 2	SE¼ sec. 26 T. 8 S., R. 5 W. Polk Co.	PTD 1,000	Application
	Reichhold Energy Corporation	Lee 32-32	NE¼ sec. 32 T. 7 N., R. 5 W. Columbia Co.	PTD 2,600	Application

(Continued on next page.)

Permit number	Operator	Well name	Location	PTD: Proposed (ft) TD: Total depth (ft)	Status
	Reichhold Energy Corporation	Paul 34-32	SE ¼ sec. 32 T. 7 N., R. 5 W. Columbia Co.	PTD 3,150	Application
	American Quasar Petroleum Co.	Benson Timber 8-14	SW ¼ sec. 8 T. 6 N., R. 4 W. Columbia Co.	PTD 5,000	Application

ABSTRACT

The Department maintains a collection of theses and dissertations on Oregon geology. From time to time we print abstracts of new acquisitions that we feel are of general interest to our readers.

GEOLOGY AND GEOCHEMISTRY OF VOLCANIC ROCKS IN THE DETROIT AREA, WESTERN CASCADE RANGE, OREGON, by Craig McKibben White (Ph.D., University of Oregon, 1980)

Data on compositional variations of igneous rocks are needed before competing models for the generation of magma at convergent plate boundaries can be realistically evaluated. The Neogene volcanic rocks of the Detroit area were selected to provide this information because they contain a relatively complete record of volcanism through most of late Cenozoic time.

The area mapped for this study is located wholly within the Western Cascade physiographic province and includes adjoining parts of the Detroit, Battle Ax, Mount Jefferson, and Breitenbush Hot Springs 15-minute quadrangles. Detailed mapping was augmented by radiometric dating in order to provide essential stratigraphic and chronological control for compositional studies.

Volcanic rocks in the Detroit area have been assigned to the following stratigraphic units: High Cascade Series (Quaternary); Outerson Formation (Pliocene); Elk Lake Formation (upper Miocene); Sardine Formation (middle Miocene); and two interfingering units, the Breitenbush Formation and the Scorpion Mountain lavas (upper Oligocene and lower Miocene). The Scorpion Mountain lavas consist of distinctive, dark-colored, generally aphanitic flows and dikes that range in composition from basaltic andesite through dacite. Flows in this unit have been mistakenly identified as Columbia River basalt; however, mapping indicates they were erupted within the Western Cascade Range, and radiometric dating has clearly shown they predate Columbia River basalt by as much as 10 m.y.

Ninety specimens from the various units exposed in the Detroit area were analyzed for ten major elements and Rb, Sr, Zr, and Ni. In addition, the abundances of Ba, Hf, Th, Cr, Sc, and seven rare-earth elements were determined for 20 samples. Compositional data show that upper Oligocene-lower Miocene lavas in the Detroit area have low contents of Al_2O_3 and high ratios of Fe/Mg characteristic of lavas in the tholeiitic rock series. Younger lavas in the map area are overwhelmingly calc-alkaline in geochemical character. Rocks in the oldest age groups also differ from younger Cascade lavas by having higher values of TiO_2 , K_2O , Rb, Zr, and Sc, and lower contents of CaO, Sr, Ni, and Cr.

The differences in major-element and included trace-element contents between the oldest lavas and those in younger suites can best be explained by a shift in the assemblage of

early-stage liquidus minerals. The trends in the oldest lavas are consistent with fractionation of olivine + plagioclase + Cr-rich spinel \pm clinopyroxene; those in the younger suites indicate fractionation of amphibole \pm olivine \pm clinopyroxene. The progressive decrease in contents of K_2O and excluded trace elements in the Detroit area lavas is compatible with a process of zone refining and depletion of mantle or lower crustal rocks in these elements with time.

Trends in the compositions of Quaternary lavas erupted from volcanic centers at Battle Ax Mountain and Mount Jefferson have been used to evaluate petrologic processes on the scale of a single volcano. Lavas in these suites show a progressive increase in SiO_2 contents with time, but data on trace-element abundances indicate each volcano produced at least three compositionally distinctive groups of flows. The initial contents of Rb, Zr, Hf, and La in each sequence of lavas are lower than those of the previously erupted group. The "resetting" of trace-element concentrations without a corresponding return to low values of SiO_2 suggests that magma reservoirs were replenished with magma that was strongly depleted in incompatible trace elements but produced lavas that were differentiated to a pre-established level. Repeated partial melting of a large volume source in the mantle could produce consecutive batches of magma with progressively lower contents of dispersed elements. □

Engineering Geologist meeting set for Portland

The National Association of Engineering Geologists will hold its twenty-fourth annual meeting this September in Portland. The meeting consists of three days of technical sessions starting on September 28, 1981. Field trips will be conducted before and after the technical sessions. Because of Portland's proximity to Mount St. Helens, special sessions and field trips are planned to focus on the engineering problems associated with the aftermath of a volcanic eruption.

Technical session topics include nuclear waste disposal; non-nuclear waste disposal; open sea engineering geology; high energy or emerging coastlines; recovery from earthquakes, volcanic events, and other geologic disasters; instrumentation; ground water; and hydrogeology. Risk analysis and liability are the topics of the symposia. Five field trips scheduled include (1) Hanford Nuclear Reservation-Columbia River, (2) Cascade Range-Central Oregon, (3) Oregon Coast, (4) Mount St. Helens-Toutle River, and (5) Mount St. Helens-Lewis River.

Details about all events appear in the January, April, and July issues of the A.E.G. Newsletter. For information or registration forms contact Ken Robbins, Dames and Moore, 1220 SW Morrison, Portland, Oregon 97205; phone (503) 228-7688.

Available publications

BULLETINS

	Price	No. Copies	Amount
33. Bibliography (1st supplement) geology and mineral resources of Oregon, 1947: Allen	\$ 1.00	_____	_____
36. Papers on Tertiary foraminifera: Cushman, Stewart and Stewart, 1949: v. 2	1.25	_____	_____
44. Bibliography (2nd supplement) geology and mineral resources of Oregon, 1953: Steere	2.00	_____	_____
46. Ferruginous bauxite deposits, Salem Hills, 1956: Corcoran and Libbey	1.25	_____	_____
49. Lode Mines, Granite mining district, Grant County, Oregon, 1959: Koch	1.00	_____	_____
53. Bibliography (3rd supplement) geology and mineral resources of Oregon, 1962: Steere and Owen	3.00	_____	_____
57. Lunar Geological Field Conference guidebook, 1965: Peterson and Groh, editors	3.50	_____	_____
62. Andesite Conference guidebook, 1968: Dole	3.50	_____	_____
65. Proceedings of the Andesite Conference, 1969: (copies)	10.00	_____	_____
67. Bibliography (4th supplement) geology and mineral resources of Oregon, 1970: Roberts	3.00	_____	_____
71. Geology of selected lava tubes in Bend area, Oregon, 1971: Greeley	2.50	_____	_____
77. Geologic field trips in northern Oregon and southern Washington, 1973	5.00	_____	_____
78. Bibliography (5th supplement) geology and mineral resources of Oregon, 1973: Roberts	3.00	_____	_____
81. Environmental geology of Lincoln County, 1973: Schlicker and others	9.00	_____	_____
82. Geologic hazards of Bull Run Watershed, Multnomah, Clackamas Counties, 1974: Beaulieu	6.50	_____	_____
83. Eocene stratigraphy of southwestern Oregon, 1974: Baldwin	4.00	_____	_____
84. Environmental geology of western Linn County, 1974: Beaulieu and others	9.00	_____	_____
85. Environmental geology of coastal Lane County, 1974: Schlicker and others	9.00	_____	_____
86. Nineteenth biennial report of the Department, 1972-1974	1.00	_____	_____
87. Environmental geology of western Coos and Douglas Counties, 1975	9.00	_____	_____
88. Geology and mineral resources of upper Chetco River drainage, 1975: Ramp	4.00	_____	_____
89. Geology and mineral resources of Deschutes County, 1976: Peterson and others	6.50	_____	_____
90. Land use geology of western Curry County, 1976: Beaulieu	9.00	_____	_____
91. Geologic hazards of parts of northern Hood River, Wasco, and Sherman Counties, Oregon, 1977: Beaulieu ..	8.00	_____	_____
92. Fossils in Oregon (reprinted from <i>The Ore Bin</i>), 1977	4.00	_____	_____
93. Geology, mineral resources, and rock material of Curry County, Oregon, 1977	7.00	_____	_____
94. Land use geology of central Jackson County, Oregon, 1977: Beaulieu	9.00	_____	_____
95. North American ophiolites, 1977	7.00	_____	_____
96. Magma genesis: AGU Chapman Conference on Partial Melting, 1977	12.50	_____	_____
97. Bibliography (6th supplement) geology and mineral resources of Oregon, 1971-75, 1978	3.00	_____	_____
98. Geologic hazards of eastern Benton County, 1979: Bela	9.00	_____	_____
99. Geologic hazards of northwestern Clackamas County, Oregon, 1979: Schlicker and Finlayson	10.00	_____	_____
100. Geology and mineral resources of Josephine County, Oregon, 1979: Ramp and Peterson	9.00	_____	_____
101. Geologic field trips in western Oregon and southwestern Washington, 1980	9.00	_____	_____

SPECIAL PAPERS

1. Mission, goals, and purposes of Oregon Department of Geology and Mineral Industries, 1978	2.00	_____	_____
2. Field geology of SW Broken Top quadrangle, Oregon, 1978: Taylor	3.50	_____	_____
3. Rock material resources of Clackamas, Columbia, Multnomah, and Washington Counties, Oregon, 1978: Gray and others	7.00	_____	_____
4. Heat flow of Oregon, 1978: Blackwell, Hull, Bowen, and Steele	3.00	_____	_____
5. Analysis and forecasts of the demand for rock materials in Oregon, 1979: Friedman and others	3.00	_____	_____
6. Geology of the La Grande area, Oregon, 1980: Barrash and others	5.00	_____	_____
7. Pluvial Fort Rock Lake, Lake County, Oregon: Allison	4.00	_____	_____
8. Geology and geochemistry of the Mt. Hood volcano, 1980: White	2.00	_____	_____
9. Geology of the Breitenbush Hot Springs quadrangle, Oregon, 1980: White	4.00	_____	_____
10. Tectonic rotation of the Oregon Western Cascades, 1980: Magill and Cox	3.00	_____	_____
12. Geologic linears of the northern part of the Cascade Range, Oregon, 1980: Venkatakrishnan, Bond, and Kauffman	3.00	_____	_____

GEOLOGIC MAPS

Geologic map of Galice quadrangle, Oregon, 1953	1.75	_____	_____
Geologic map of Albany quadrangle, Oregon, 1953	1.00	_____	_____
Reconnaissance geologic map of Lebanon quadrangle, 1956	1.50	_____	_____
Geologic map of Bend quadrangle and portion of High Cascade Mountains, 1957	1.50	_____	_____
Geologic map of Oregon west of 121st meridian (USGS I-325), 1961	2.25	_____	_____
Geologic map of Oregon east of 121st meridian (USGS I-902), 1977	3.75	_____	_____
GMS-4: Oregon gravity maps, onshore and offshore, 1967 (folded)	3.00	_____	_____
GMS-5: Geologic map of Powers quadrangle, Oregon, 1971	2.00	_____	_____
GMS-6: Preliminary report on geology of part of Snake River Canyon, 1974	6.50	_____	_____
GMS-7: Geology of the Oregon part of the Baker quadrangle, Oregon, 1976	3.00	_____	_____
GMS-8: Complete Bouguer gravity anomaly map, Cascade Mountain Range, central Oregon, 1978	3.00	_____	_____
GMS-9: Total field aeromagnetic anomaly map, Cascade Mountain Range, central Oregon, 1978	3.00	_____	_____
GMS-10: Low- to intermediate-temperature thermal springs and wells in Oregon, 1978	2.50	_____	_____
GMS-12: Geologic map of the Oregon part of the Mineral quadrangle, 1978	2.00	_____	_____
GMS-13: Geologic map of the Huntington and part of the Olds Ferry quadrangles, Oregon, 1979	3.00	_____	_____
GMS-15: Free-air gravity anomaly map and complete Bouguer gravity anomaly map, Cascade Mountain Range, northern Oregon, 1981	3.00	_____	_____
GMS-16: Free-air gravity anomaly map and complete Bouguer gravity anomaly map, Cascade Mountain Range, southern Oregon, 1981	3.00	_____	_____
GMS-17: Total-field aeromagnetic anomaly map, Cascade Mountain Range, southern Oregon, 1981	3.00	_____	_____

SHORT PAPERS

18. Radioactive minerals the prospector should know, 1976: White, Schafer, Peterson75	_____	_____
21. Lightweight aggregate industry in Oregon, 1951: Mason25	_____	_____
24. The Almeda Mine, Josephine County, Oregon, 1967: Libbey	3.00	_____	_____
25. Petrography, type Rattlesnake Formation, central Oregon, 1976: Enlows	2.00	_____	_____
27. Rock material resources of Benton County, 1978: Schlicker and others	4.00	_____	_____

Available publications (continued)

MISCELLANEOUS PAPERS

	Price	No. Copies	Amount
1. A description of some Oregon rocks and minerals, 1950: Dole	\$ 1.00	_____	_____
5. Oregon's gold placers (reprints), 195450	_____	_____
8. Available well records of oil and gas exploration in Oregon, rev. 1980: Olmstead and Newton	2.00	_____	_____
11. Collection of articles on meteorites, 1968 (reprints from <i>The Ore Bin</i>)	1.50	_____	_____
13. Index to <i>The Ore Bin</i> , 1950-1974	1.50	_____	_____
14. Thermal springs and wells, 1970: Bowen and Peterson (with 1975 supplement)	1.50	_____	_____
15. Quicksilver deposits in Oregon, 1971: Brooks	1.50	_____	_____
17. Geologic hazards inventory of the Oregon coastal zone, 1974: Beaulieu, Hughes, and Mathiot	5.00	_____	_____
18. Proceedings of Citizens' Forum on potential future sources of energy, 1975	2.00	_____	_____
19. Geothermal exploration studies in Oregon—1976, 1977	3.00	_____	_____
20. Investigations of nickel in Oregon, 1978: Ramp	5.00	_____	_____

OIL AND GAS INVESTIGATIONS

3. Preliminary identifications of foraminifera, General Petroleum Long Bell #1 well	2.00	_____	_____
4. Preliminary identifications of foraminifera, E.M. Warren Coos County 1-7 well, 1973	2.00	_____	_____
5. Prospects for natural gas production or underground storage of pipeline gas, upper Nehalem River Basin, Columbia-Clatsop Counties, Oregon, 1976	5.00	_____	_____
6. Prospects for oil and gas in the Coos Basin, western Coos, Douglas, and Lane Counties, Oregon, 1980: Newton and others	9.00	_____	_____

MISCELLANEOUS PUBLICATIONS

Landforms of Oregon (17 × 12 inches)50	_____	_____
Mining claims (State laws governing quartz and placer claims)50	_____	_____
Geological highway map, Pacific NW region, Oregon-Washington (published by AAPG)	5.00	_____	_____
Fifth Gold and Money Session and Gold Technical Session Proceedings, 1975	5.00	_____	_____
Sixth Gold and Money Session and Gold Technical Session Proceedings, 1978	6.50	_____	_____
Back issues of <i>The Ore Bin</i>	25¢ over the counter; 75¢ mailed	_____	_____
Back issues of <i>Oregon Geology</i>	40¢ over the counter; 75¢ mailed	_____	_____
Colored postcard, <i>Geology of Oregon</i>	10¢ each; 3 for 25¢; 7 for 50¢; 15 for 1.00	_____	_____

OREGON GEOLOGY

1005 State Office Building, Portland, Oregon 97201

Second Class Matter

POSTMASTER: Form 3579 requested

PUBLICATIONS ORDER

Omission of price indicates publication is in press. Minimum mail order 50¢. All sales are final. Publications are sent postpaid. Payment must accompany orders of less than \$20.00.

Fill in appropriate blanks and send sheet to Department (see address on reverse side).

YOUR NAME _____

ADDRESS _____

Zip _____

Amount enclosed \$ _____

OREGON GEOLOGY

____ Renewal ____ Subscription ____ Gift

____ 1 Year (\$4.00) ____ 3 Years (\$10.00)

NAME _____

ADDRESS _____

____ ZIP _____

(If Gift, From: _____)