



The Oregon Department of Geology and Mineral Industries (DOGAMI) has been identifying and mapping the tsunami inundation hazard along the Oregon coast since 1994. In Oregon, DOGAMI manages the National Tsunami Hazard Mitigation Program, which has been administered by the National Oceanic and Atmospheric Administration (NOAA) since 1995. DOGAMI's work is designed to help cities, counties, and other states in coastal areas reduce the potential for disastrous tsunami-related consequences by understanding and mitigating this geologic hazard. Using federal funding awarded by NOAA, DOGAMI has developed a new generation of tsunami inundation maps to help residents and visitors along the entire Oregon coast prepare for the next Cascadia Subduction Zone (CSZ) earthquake and tsunami.

The CSZ is the tectonic plate boundary between the North American Plate and the Juan de Fuca Plate (Figure 1). These plates are converging at a rate of about 1.5 inches per year, but the movement is not smooth and continuous. Rather, the plates lock in place, and unreleased energy builds over time. At intervals, this accumulated energy is violently released in the form of a megathrust earthquake rupture, where the North American Plate suddenly slips westward over the Juan de Fuca Plate. This rupture causes a vertical displacement of water that creates a tsunami (Figure 2). Similar rupture processes and tsunamis have occurred elsewhere on the planet where subduction zones exist: for example, offshore Chile in 1950 and 2010, offshore Alaska in 1954, near Sumatra in 2004 and offshore Japan in March 2011.

Oregon, DOGAMI has also incorporated physical evidence that suggests that portions of the coast may drop 4 to 10 feet during the earthquake; this effect is known as subsidence. Detailed information on fault geometries, subsidence, computer models, and the methodology used to create the tsunami scenarios presented on this map can be found in DOGAMI Special Papers 41 (Priest and others, 2009) and 43 (Witter and others, 2011).

This tsunami inundation map displays the output of computer models representing five selected tsunami scenarios, all of which include the earthquake-produced subsidence and the tsunami-amplifying effects of the spillaway fault. (MHHW stands for a tsunami occurs at Mean Higher High Water (MHHW) tide; MHHW is defined as the average height of the higher high tides observed over an 18-year period at the Yaquina Bay (Central Coast Model) tide gauge. To make it easier to understand this scientific material and to enhance the educational aspects of hazard mitigation and response, the five scenarios are labeled as "T-shirt sizes" ranging from Small, Medium, Large, Extra Large, to Extra Extra Large (S, M, L, XL, XXL). The map legend depicts the respective amounts of slip, the frequency of occurrences, and the earthquake magnitude for these five scenarios. Figure 4 shows the cumulative number of buildings inundated within the map area.

The computer simulation model output is provided to DOGAMI as millions of points with values that indicate whether the location of each point is wet or dry. These points are converted to wet and dry contour lines that form the extent of inundation. The transition area between the wet and dry contour lines is termed the Wet/Dry Zone, which equates to the amount of error in the model when determining the maximum inundation for each scenario. Only the XGL Wet/Dry Zone is shown on this map.

CSZ Model Specifications: The sizes of the earthquake and its resultant tsunami are primarily driven by the amount and geometry of the slip that takes place when the North American Plate snags westward over the Juan de Fuca Plate during a CSZ event. DOGAMI has modeled a wide range of earthquake and tsunami sizes that take into account different fault geometries that could amplify the amount of seafloor displacement and increase tsunami inundation. Seismic geophysical profiles show that there may be a steep splay fault running nearly parallel to the CSZ but closer to the Oregon coastline (Figure 3). The effect of this splay fault moving during a full-rupture CSZ event would be an increase in the amount of vertical displacement of the Pacific Ocean, resulting in an increase of the tsunami inundation onshore in

A cross-sectional diagram of the Juan de Fuca Plate subducting beneath the North American Plate. The Pacific Plate is shown on the left, with the Juan de Fuca Plate extending from it. The Juan de Fuca Plate is shown subducting under the North American Plate. The subducting plate is labeled 'Juan de Fuca Plate' and the overriding plate is labeled 'NORTH AMERICAN PLATE'. The boundary between them is labeled 'JUAN DE FUCA TRANSFORM' and 'JUAN DE FUCA SUBDUCTION ZONE'. The subducting plate is shown dipping into the mantle, with a 'Locked Zone' indicated. The mantle is labeled 'Magma' at two points. The diagram also shows the 'Cascadia Subduction Zone' and 'Washington Divergent'.

Figure 1: This block diagram depicts the tectonic setting of the region. See Figure 2 for the sequence of events that occur during a Cascadia Subduction Zone megathrust earthquake and tsunami.

A

Figure 2: The North American Plate rides over the descending Juan de Fuca Plate at a rate of approximately 1.5 inches per year.

B

Because the two plates are stuck in place at the "locked zone," strain builds up over time and the North American Plate bulges up.

C

Eventually, the locked zone ruptures and causes a great earthquake. The sudden slip of the two plates displaces Pacific Ocean water upward and creates a tsunami.

D

Displaced and uplifted Pacific Ocean water rushes in all directions.

E

Along the Oregon coast, tsunami waves run up onto the beach for several hours.

Figure 3 is a bar chart titled "Timing, frequency, and magnitude of the last 19 great Cascadia Subduction Zone events over the past 3000 years". The x-axis is labeled "Research-indicated radiocarbon age of last event [most recent in January 1700]" and ranges from 3000 to 0 years. The y-axis is labeled "Average Offshore Landslide Turbidity Magnitude" and ranges from 0 to 10. The chart displays 19 events, each represented by a bar. The bars are color-coded: red for "larger but less frequent turbidities" (labeled XXL, XL, L) and blue for "smaller but more frequent turbidities" (labeled M, S). The red bars are generally taller, indicating higher turbidity magnitudes, while the blue bars are shorter, indicating lower magnitudes. The red bars are clustered between 1700 and 3000 years, with the tallest bar (XXL) occurring just before 1700 AD. The blue bars are clustered between 0 and 1700 AD, with the shortest bar (S) occurring just after 1700 AD. A vertical line at 1700 AD marks the present time. The legend indicates that the red bars are "larger but less frequent turbidities" and the blue bars are "smaller but more frequent turbidities".

	Entire Map Area	City of Landside	Unincorporated
Total Buildings	2,101	1,603	498
Buildings within Tsunami Zones*			
Small	1	0	1
Medium	1	0	29
Large	151	4	147
Extra Large	191	38	153
Extra Extra Large	231	75	156
Percent of Buildings within Tsunami Zones			
Small	0.0%	0.0%	0.2%
Medium	1.4%	0.0%	5.8%
Large	7.2%	0.2%	29.5%
Extra Large	9.1%	2.4%	30.7%
Extra Extra Large	11.0%	4.7%	31.3%

*Building counts shown are based on polygon centroids and are cumulative within the map area.

*Building counts shown are based on polygon centroids and are cumulative within the map area.




Figure 4: The table and chart show the number of buildings inundated for each "tsunami T-shirt scenario" for cities and unincorporated portions of the map.

Line graph showing Elevation, feet (NAVD83) on the Y-axis (0.0 to 50.0) versus Time since Earthquake (hours) on the X-axis (0.0 to 8.0). The graph displays the change in water level for five tsunami scenarios (Xxl, xl, L, M, S) at the simulated gauge station shown on the map (Tenmile Creek (Station C5)).


Legend:

- Xxl: Change in water level for five tsunami scenarios
- xl: Change in water level for five tsunami scenarios
- L: Change in water level for five tsunami scenarios
- M: Change in water level for five tsunami scenarios
- S: Change in water level for five tsunami scenarios

Station: Tenmile Creek (Station C5)

Figure 5: This chart depicts the tsunami waves as they arrive at the selected reference point (simulated gauge station). It shows the change in wave heights for all five tsunami scenarios over an 8-hour period. The starting wave elevation (0.0 hours) takes into account the local land subsidence or uplift caused by the earthquake. Wave heights vary through time, and the first wave will not necessarily be the largest as waves interfere and reflect off local topography and bathymetry. Any absence of data indicates periods for which tsunami inundation has not yet reached or has receded from the station location and dry land is exposed.

Figure 6: These profiles depict the expected maximum tsunami wave elevation for the five "tsunami T-shirt scenarios" along lines A-A' and B-B'. The tsunami scenarios are modeled to occur at high tide and to account for local subsidence or uplift of the ground surface.



Earthquake size	Average Slip Range (ft)	Maximum Slip Range (ft)	Time to Accumulate Slip (yrs)	Earthquake Magnitude
	XXL 59 to 72	118 to 144	1,200	-9.1
	XL 56 to 72	115 to 144	1,050 to 1,200	-9.1
	L 36 to 49	72 to 98	650 to 800	-9.0
	M 23 to 30	46 to 62	425 to 525	-8.9
	S 13 to 16	30 to 36	300	-8.7
	XXL Wet/Dry Zone			

Coos-01 Lakeside West
Coos-02 Lakeside East
Coos-10 Ilwaco Slough
Coos-11 Catching Slough

[illegible]

2012

**Tsunami Inundation Maps for Lakeside West,
Coos County, Oregon**

Plate 1

